FORM PTO	0-1390 ILS DEPARTMENT OF	COMMERCE PATENT AND TRADEMARK OFFICE	1
(REV. 11-2	2000)	ER TO THE UNITED STATES	ATTORNEY'S DOCKET NUMBER 58049-00002
		CTED OFFICE (DO/EO/US)	US APPLICATION NO (If known, sec 37 CFR 15
	CONCERNING A FIL	ING UNDER 35 U.S.C. 371	10/088548
	NATIONAL APPLICATION NO		PRIORITY DATE CLAIMED
	CR99/00554 OF INVENTION	September 15, 1999	September 15, 1999
A NO	VEL ANGIOGENESIS INHIBIT	OR .	
CHAN		RK, Eun Jeong; YUM, Jung-sun; CHUNG, Soo	
Applica	int herewith submits to the United	States Designated/Elected Office (DO/EO/US)	the following items and other information:
_	•	ems concerning a filing under 35 U.S.C. 371.	
2.		JENT submission of items concerning a filing t	
3.	items (5), (6), (9) and (21) indicate		
4.		xpiration of 19 months from the priority date (A	article 31).
3.		cation as filed (35 U.S C. 371(c)(2))	
	· <u></u>	ired only if not communicated by the Internation	nal Bureau).
		by the International Bureau.	
ku -		oplication was filed in the United States Receive	- ,
6.		of the International Application as filed (35 U.S.	C. 371(c)(2)).
	a. is attached hereto.		
_		omitted under 35 U.S.C. 154(d)(4).	
7. 🔲		International Aplication under PCT Article 19 (
	a. are attached hereto (req	uired only if not communicated by the Internation	onal Bureau).
	b. have been communicate	ed by the International Bureau.	
		wever, the time limit for making such amendme	ents has NOT expired.
	d. have not been made and		
· <u> </u>		f the amendments to the claims under PCT Artic	cle 19 (35 U.S.C. 371 (c)(3)).
9. 🗸	An oath or declaration of the inve	ntor(s) (35 U.S.C. 371(c)(4)).	-
10.	An English lanugage translation of Article 36 (35 U.S.C. 371(c)(5)).	f the annexes of the International Preliminary È	xamination Report under PCT
Item	s 11 to 20 below concern docum	ent(s) or information included:	
н. 🔲		ement under 37 CFR 1.97 and 1.98.	
12.	An assignment document for re	cording A separate cover sheet in compliance	with 37 CFR 3.28 and 3.31 is included.
13.	A FIRST preliminary amendme		
14.	A SECOND or SUBSEQUENT	preliminary amendment	
15.	A substitute specification.		
16.	A change of power of attorney	and/or address letter.	
17. 🗸	A computer-readable form of th	e sequence listing in accordance with PCT Rule	13ter.2 and 35 U S.C. 1.821 - 1.825.
18.	A second copy of the published	international application under 35 U.S.C. 154(d	I)(4)
19.	A second copy of the English la	nguage translation of the international application	on under 35 U S C 154(d)(4).
20. 🗸	Other items or information: M	icroorganism Deposit Rec earch Report	eipt; International
		`	

U.S. APPLITATION NO (1 C)	88548	INTERNATIONAL APPLICATION NO			ATTORNEY'S DOC	KET NUMBER						
21. The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)): Neither international preliminary examination fee (37 CFR 1 482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO. \$1000.00												
International prelin USPTO but Intern	ninary examination fee o ational Search Report p											
	ninary examination fee (earch fee (37 CFR 1.445											
but all claims did n	ot satisfy provisions of	(37 CFR 1.482) paid to US PCT Article 33(1)-(4)	\$690.00									
International prelin	ninary examination fee (fied provisions of PCT .	(37 CFR 1.482) paid to US Article 33(1)-(4)	SPTO \$100.00		· · · · · · · · · · · · · · · · · · ·							
ENTE	R APPROPRIATE	BASIC FEE AMO	UNT =	\$ 1,0	00.00							
	0 for furnishing the oath liest claimed priority da	\$										
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$								
Total claims	19 - 20 =	0	x \$18.00	\$	10.00							
Independent claims	$\begin{array}{ccc} & 6 & -3 = \\ \hline DENT CLAIM(S) & \text{(if approximate)} \end{array}$		x \$80.00	\$ 2 \$	40.00							
MOETIFEE DEFEN		·	+ \$270.00		240.00							
Applicant claim are reduced by	s small entity status. Se	ce 37 CFR 1.27. The fees			240.00 520.00							
		SI	UBTOTAL =	\$ 6	520.00							
Processing fee of \$1 months from the earl	30.00 for furnishing the liest claimed priority da	English translation later the		\$		-						
		TOTAL NATIO	NAL FEE =	\$ 6	20.00							
Fee for recording the accompanied by an a	e enclosed assignment (2 appropriate cover sheet (37 CFR 1.21(h)). The assi (37 CFR 3.28, 3.31). \$40.	gnment must be 00 per property +	\$	40.00							
		TOTAL FEES E	NCLOSED =	\$ (560.00	스탠드 V						
					unt to be efunded:	\$						
					charged:	\$						
a. 🗸 A check in	the amount of \$	660.00 to cover the	e above fees is enclos	sed								
b. Please char A duplicate	ge my Deposit Account copy of this sheet is en	No. in closed.	the amount of \$		to cover the	above fees.						
		rized to charge any addition of the charge any addition of the charge and addition of the charge and the charge				ny						
d. Fees are to linformation	be charged to a credit can should not be include	rd. WARNING: Informed on this form. Provide	ation on this form mag	y becon	me public. Creature	dit card PTO-2038.						
		under 37 CFR 1.494 or 1 to restore the application		et, a p	etition to reviv	e (37 CFR						
801 South Figure Los Angeles, (Tel: (213) 624 Fax: (213) 623	rs & Dempsey L.L ueroa Street, 14th California 90017-5 4-2500 3-4581	Floor 554	SIGNATU		pul L	Ph D.						
Attn: Joseph I	Hyosuk Kim, Ph.D		REGISTRA	MION	NUMBER ,	41,425						

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Jihoon Chang et al.) Group Art Unit: Not Yet Determined										
Serial No.: Not Assigned Yet) Examiner: Not Yet Determined										
Based On PCT/KR99/00554 filed on Sep. 15,)·)·										
1999 Filed: March 15, 2002)										
For: A NOVEL ANGIOGENESIS INHIBITOR)))										
PRELIMINARY AMENDMENT, S	SUBMISSION OF SEQUENCE LISTING, AND										
SUBMISSION OF MICROORGANISM DEPOSIT RECEIPT											
BOX SEQ Commissioner for Patents Washington, D.C. 20231											
Sir:	N N										
Prior to examination of the above-iden	ntified application on the merits, please amend the										
application as follows:											
	TE OF MAILING F.R. § 1.10)										
deposited with the United States Postal Service	referred to as being attached or enclosed) is being e on the date shown below with sufficient postage an envelope addressed to Box Patent Application, 20231.										
EL 824098099 US <u>M</u>	Margie A. Uribe										
Express Mail Label No.	ame of Person Mailing Paper										
March 15, 2002	// /arde/ /rube										
Date of Deposit LosAngeles/75109.1	ignature of Person Mailing Paper										

Express Mail Label No. EL 824098099 US

10088107088548

PATENT 58049-00002 JC10 Rec'd PCT/PTO 1 5 MAR 2002

IN THE SPECIFICATION:

At page 1, line 3, add the following:

-- CROSS REFERENCE TO OTHER APPLICATIONS

The present application is filed under 35 U.S.C. 371, and is the U.S. national phase application of PCT/KR99/00554, filed on September 15, 1999. --

IN THE CLAIMS:

Please amend claims as follows:

- 1. (Amended) LK6 protein comprising amino acid sequences of human apolipoprotein(a) kringle domains IV36.
- 2. (Amended) LK7 protein comprising amino acid sequences of human apolipoprotein(a) kringle domains IV37.
- 3. (Amended) LK8 protein comprising amino acid sequences of human apolipoprotein(a) kringle domains V38.
- 4. (Amended) LK68 protein comprising amino acid sequences of human apolipoprotein(a) kringle domains IV36, IV37 and V38 in a serial manner
 - 5. (Amended) A cDNA sequence which codes for the LK6 protein of claim 1.

- 6. (Amended) A cDNA sequence which codes for the LK7 protein of claim 2.
- 7. (Amended) A cDNA sequence which codes for the LK8 protein of claim 3.
- 8. (Amended) A cDNA sequence which codes for the LK68 protein of claim 4.

REMARKS

Claims 1-19 are pending in the application. The amendments to the claims serve to further clarify the present invention. No new matter has been inserted into the application. Accordingly, entry of the amendments to the application and examination on the merits are respectfully requested.

Submission Of Sequence Listing

Enclosed herewith in full compliance with 37 C.F.R. §§ 1.821-1.825 is a Sequence Listing to be inserted into the specification. The Sequence Listing in no way introduces new matter into the specification.

Also submitted herewith in full compliance with 37 C.F.R. §§ 1.821-1.825 is a computer readable disk copy of the Sequence Listing. Applicant's undersigned representative hereby states that the information recorded in computer readable form is identical to the written sequence listing.

Submission of Microorganism Deposit Receipt

Applicants submit herewith the deposit receipts for *Escherichia coli* BL21/LK6-8 (KCTC 0633BP) and *Escherichia coli* BL21/LK8 (KCTC 0634BP) which were deposited with the Korean Collection for Type Cultures on June 9, 1999, and *Escherichia coli* BL21(DE3)/LK6 (KCTC

achersell of aspir

Express Mail Label No. EL 824098099 US

PATENT 58049-00002

0655BP) and *Escherichia coli* BL21(DE3)/LK7 (KCTC 0656BP) which were deposited with the Korean Collection for Type Cultures on Sept. 3, 1999. Applicants hereby declare that subject to paragraph (b) of 37 CFR 1.808, all restrictions imposed by the depositor on the availability to the public of the deposited material will be irrevocably removed upon the granting of the patent.

The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 07-1853 during the pendency of prosecution of this application. A duplicate of this paper is enclosed for the Deposit Account, should it be needed.

Respectfully submitted,

SQUIRE, SANDERS & DEMPSEY L.L.P.

Dated: March 15, 2002

seph Hyosuk Kim, Ph.D.

Reg. No. 41.425

801 S. Figueroa Street, 14th Floor Los Angeles, CA 90017-5554 Telephone: (213) 689-6533

Facsimile: (213) 623-4581

VERSION MARKED TO SHOW CHANGES MADE

In the Specification

At page 1, line 3, add the following:

-- CROSS REFERENCE TO OTHER APPLICATIONS

The present application is filed under 35 U.S.C. 371, and is the U.S. national phase application of PCT/KR99/00554, filed on September 15, 1999. --

In the Claims

Please amend claims 1-8 as follows:

- 1. (Amended) LK6 protein [(SEQ ID NO: 4) consisting of] <u>comprising</u> amino acid sequences of human apolipoprotein(a) kringle domains IV36.
- 2. (Amended) LK7 protein [(SEQ ID NO: 6) consisting of] <u>comprising</u> amino acid sequences of human apolipoprotein(a) kringle domains IV37.
- 3. (Amended) LK8 protein [(SEQ ID NO: 8) consisting of] <u>comprising</u> amino acid sequences of human apolipoprotein(a) kringle domains V38.
- 4. (Amended) LK68 protein [(SEQ ID NO: 2) consisting of] comprising amino acid sequences of human apolipoprotein(a) kringle domains IV36, IV37 and V38 in a serial manner

- 5. (Amended) A cDNA sequence [(SEQ ID NO: 3)] which codes for the LK6 protein of claim 1.
- 6. (Amended) A cDNA sequence [(SEQ ID NO: 5)] which codes for the LK7 protein of claim 2.
- 7. (Amended) A cDNA sequence [(SEQ ID NO: 7)] which codes for the LK8 protein of claim 3.
- 8. (Amended) A cDNA sequence [(SEQ ID NO: 1)] which codes for the LK68 protein of claim 4.

10/088548 JC10 Rec'd PCT/PTO 1 5 MAR 2002

SEQUENCE LISTING

,	<110>	Mogam Biotechnology Research	Institute et al	
5	<120>	A NOVEL ANGIOGENESIS INHIBITOR	?	
	<160>	14		
10	<170>	KOPATIN 1.5		
	<210>	1		
	<211>	924		
	<212>	DNA		
15	<213>	Homo sapiens		
	<400>	1		
	aaaagccci	g tggtccagga ttgctaccat ggtgat	ggac ggagttatcg aggcatatcc	60
			gone gong to be a magenture of	00
20	tccaccact	g tcacaggaag gacctgtcaa tcttgg	tcat ctatgatacc acactggcat	120
	cagaggaco	c cagaaaacta cccaaatgct ggcctg	accg agaactactg caggaatcca	180
\ -	gattctggg	a aacaaccctg gtgttacaca accgat	ccgt gtgtgaggtg ggagtactgc	240
25	aatctgaca	c aatgeteaga aacagaatea ggtgte	ctag agactcccac tgttgttcca	300
	gttccaago	a tggaggetea ttetgaagea geacea	actg agcaaacccc tgtggtccgc	360
30	cagtgctac	c atggcaatgg ccagagttat cgaggc	acat tetecaceae tgteacagga	420
	aggacatgt	c aatetiggie atecatgaca ecacae	cggc atcagaggac cccagaaaac	480
	tacccaaat	g atggeetgae aatgaactae tgeagg	aate cagatgeega tacaggeeet	540
5	tggtgtttt	a ccacggaccc cagcatcagg tgggagi	tact gcaacctgac gcgatgctca	600

	gacaca	agaag	ggao	etgtg	gto	gctc	ctcc	g ac	tgtc	atco	agg	ttco	aag	ccta	igggco	t	660
	ccttci	gaac	aaga	ectgt	at g	gtttg	ggaa	t gg	gaaa	ıggat	acc	gggg	gcaa	gaag	gcaac	cc	720
5	actgtt	actg	ggao	gcca	tg c	cagg	aatg	g go	tgcc	cagg	ago	ccca	ıtag	acac	agcac	g .	780
	ttcatt	ccag	ggac	aaat	aa a	ıtggg	cagg	t ct	ggaa	aaaa	att	actg	ccg	taac	cctga	it	840
1.0	ggtgad	atca	atgg	tccc	tg g	tgct	acac	a at	gaat	ccaa	gaa	aact	ttt	tgac	tactg	;t	900
10	gatato	cctc	tctg	tgca	tc c	tct											924
	<210>	2														•	
15	<211>	30)8														
	<212>	PF															
	<213>		omo s	apie	ns												
	<400>	2															
20	Lys Se	r Pro	Val	Val 5	Gln	Asp	Cys	Tyr	His 10	Gly	Asp	Gly	Arg	Ser 15			
	•			J					10					13			
	Arg G1	y Ile	Ser 20	Ser	Thr	Thr	Val	Thr 25	Gly	Arg	Thr	Cys	G1n 30		Trp		
25													00				
	Ser Se	r Met 35		Pro	His	Trp	His 40	Gln	Arg	Thr	Pro	Glu 45	Asn	Tyr	Pro		
30	Asn Al		Leu	Thr	Glu	Asn 55		Cys	Arg	Asn	Pro 60		Ser	Gly	Lys		
	Gln Pr	o Trp	Cys	Tyr	Thr 70	Thr	Asp	Pro	Cys	Val 75	Arg	Trp	Glu	Tyr	Cys 80		
35	Asn Le	u Thr	Gln	Cys 85	Ser	Glu	Thr	Glu	Ser	Gly	Val	Leu	Glu	Thr	Pro		

	Thr	Val	Val	Pro 100		Pro	Ser	Met	G1u 105		His	Ser	Glu	Ala 110		Pro
5	Thr	Glu	Gln 115		Pro	Val	Val	Arg 120		Cys	Tyr	His	Gly 125		Gly	Gln
	Ser	Tyr 130		Gly	Thr	Phe	Ser 135	Thr	Thr	Val	Thr	Gly 140		Thr	Cys	Gln
10	Ser 145		Ser	Ser	Met	Thr 150	Pro	His	Arg	His	Gln 155	Arg	Thr	Pro	Glu	Asn 160
15	Tyr	Pro	Asn	Asp	Gly 165	Leu	Thr	Met	Asn	Tyr 170	Cys	Arg	Asn	Pro	Asp 175	Ala
15	Asp	Thr	Gly	Pro 180	Trp	Cys	Phe	Thr	Thr 185	Asp	Pro	Ser	He	Arg 190	Trp	Glu
20	Tyr	Cys	Asn 195	Leu	Thr	Arg	Cys	Ser 200	Asp	Thr	Glu	Gly	Thr 205	Val	Val	Ala
	Pro	Pro 210	Thr	Val	He	Gln	Val 215	Pro	Ser	Leu	Gly	Pro 220	Pro	Ser	Glu	Gln
25	Asp 225	Cys	Met	Phe	Gly	Asn 230	Gly	Lys	Gly	Tyr	Arg 235	Gly	Lys	Lys	Ala	Thr 240
	Thr	Val	Thr	Gly	Thr 245	Pro	Cys	Gln	Glu	Trp 250	Ala	Ala	Gln	Glu	Pro 255	His
30	Arg	His	Ser	Thr 260	Phe	He	Pro	Gly	Thr 265	Asn	Lys	Trp	Ala	Gly 270	Leu	Glu
35	Lys	Asn	Tyr 275	Cys	Arg	Asn	Pro	Asp 280	Gly	Asp	He	Asn	Gly 285	Pro	Trp	Cys
	Tyr	Thr	Met	Asn	Pro	Arg	Lys	Leu	Phe	Asp	Tyr	Cys	Asd	Ile	Pro	Leu

290 295 300

Cys Ala Ser Ser
305

5

<400>

aaaagccctg tggtccagga ttgctaccat ggtgatggac ggagttatcg aggcatatcc 60

tccaccactg tcacaggaag gacctgtcaa tcttggtcat ctatgatacc acactggcat 120

cagaggaccc cagaaaacta cccaaatgct ggcctgaccg agaactactg caggaatcca 180

gattetggga aacaaccetg gtgttacaca accgatecgt gtgtgaggtg ggagtactgc 240

aatetgacac aatgeteaga aacagaatea ggt 273

<210> 4
25 <211> 91
<212> PRT
<213> Homo sapiens

35

<400> 4
Lys Ser Pro Val Val Gln Asp Cys Tyr His Gly Asp Gly Arg Ser Tyr
1 5 10 15

Arg Gly Ile Ser Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ser Trp
20 25 30

Ser Ser Met Ile Pro His Trp His Gln Arg Thr Pro Glu Asn Tyr Pro 35 40 45

	Asn Ala Gly Leu Thr Glu Asn Tyr Cys Arg Asn Pro Asp Ser Gly Lys 50 55 60	
5	Gln Pro Trp Cys Tyr Thr Thr Asp Pro Cys Val Arg Trp Glu Tyr Cys 65 70 75 80	
10	Asn Leu Thr Gln Cys Ser Glu Thr Glu Ser Gly 85 90	
15	<210> 5 <211> 267 <212> DNA <213> Homo sapiens	
	<400> 5 gtccgccagt gctaccatgg caatggccag agttatcgag gcacattctc caccactgtc	60
20	acaggaagga catgtcaatc ttggtcatcc atgacaccac accggcatca gaggacccca	120
	gaaaactacc caaatgatgg cctgacaatg aactactgca ggaatccaga tgccgataca	180
25	ggcccttggt gttttaccac ggaccccagc atcaggtggg agtactgcaa cctgacgcga	240
	tgctcagaca cagaagggac tgtggtc	267
30	<210> 6 <211> 89 <212> PRT <213> Homo sapiens	
15	<pre><400> 6 Val Arg Gln Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly Thr Phe 1 5 10 15</pre>	

	Ser Thr	Thr Val	Thr Gly	Arg Thr	Cys Gln 25	Ser Trp	Ser Ser 30		
5	Pro His	Arg His 35	Gln Arg	Thr Pro	Glu Asn	Tyr Pro	Asn Asp 45	Gly Leu	
	Thr Met	Asn Tyr	Cys Arg	Asn Pro 55	Asp Ala	Asp Thr		Trp Cys	
10	Phe Thr	Thr Asp	Pro Ser 70	Ile Arg	Trp Glu	Tyr Cys 75	Asn Leu	Thr Arg	
,	Cys Ser	Asp Thr	Glu Gly 85	Thr Val	Val				,
15									
	<210>	7							
	<211>	258							
	<212>	DNA							
20	<213>	Homo sa	piens						
	<400>	7							
	gaacagga	ct gcatg	tttgg ga	atgggaaa	ggatacc	ggg gcaa	ngaaggc a	accactgtt	60
25	actgggac	gc catgc	cagga at	gggctgcc	caggage	ccc atag	gacacag c	cacgttcatt	120
	ccagggac	aa ataaa	tgggc ag	gtctggaa	aaaaatt	act gccg	gtaaccc t	gatggtgac	180
	atcaatgg	tc cctgg	tgcta ca	caatgaat	ccaagaa	aac tttt	tgacta c	tgtgatatc	240
30	cctctctg	tg catcc	tct						258
	<210>	8							
35	<211>	86							
	<212>	PRT							
	<213>	Homo sa	oiens						
		1	-						

<400> 8

Glu Gln Asp Cys Met Phe Gly Asn Gly Lys Gly Tyr Arg Gly Lys Lys

1

5

10

15

5

Ala Thr Thr Val Thr Gly Thr Pro Cys Gln Glu Trp Ala Ala Gln Glu
20 25 30

Pro His Arg His Ser Thr Phe Ile Pro Gly Thr Asn Lys Trp Ala Gly

10 35

40

45

Leu Glu Lys Asn Tyr Cys Arg Asn Pro Asp Gly Asp Ile Asn Gly Pro 50 55 60

Trp Cys Tyr Thr Met Asn Pro Arg Lys Leu Phe Asp Tyr Cys Asp IIe
65 70 75 80

Pro Leu Cys Ala Ser Ser

85

20

<210> 9

<211> 29

<212> DNA

25 <213> Artificial Sequence

<220>

<223> single standed oligonucleotide

30 <400> 9

tccatatgaa aagccctgtg gtccaggat

29

<210> 10

35 <211> 33

<212> DNA

<213> Artificial Sequence

31

26

<220> <223> single stranded oligonucleotide <400> 10 cagtecatat ggteegeeag tgetaceatg gea <210> 11 <211> 31 10 <212> DNA <213> Artificial Sequence <220> <223> 15 single stranded olgonucleotide <400> 11 ggaattccat atggaacagg actgcatgtt t 20 <210> 12 <211> 26 <212> DNA <213> Artificial Sequence 25 <220> <223> single stranded oligonucleotide <400> 12 cgggatcctt aacctgattc tgtttc 30 <210> 13 <211> 26 <212> DNA 35 <213> Artificial Sequence

<220>

<223> single stranded oligonucleotide

<400> 13

5 cgggateett agaccaeagt eeette

26

<210> 14

<211> 23

10 <212> DNA

<213> Artificial Sequence

<220>

<223> single stranded oligonucleotide

15

<400> 14

cgggatcctt aagaggatgc aca

23

WO 01/19868

12/1/213 1

10/088548

A NOVEL ANGIOGENESIS INHIBITOR PCT/PTO 1 5 MAR 2002

BACKGROUND OF THE INVENTION

Field of the Invention

invention present relates to angiogenesis inhibitor, LK68 whose amino acid sequence is identical with the human apolipoprotein(a) kringle domains IV36, IV37 and V38, more specifically, to an acid sequence of the LK68, a cDNA sequence encoding the LK68, a recombinant expression vector comprising the CDNA, а recombinant microorganism transformed with the recombinant expression vector and a novel use of the LK68 as an anticancer agent and a method for treating the angiogenesis-mediated disease.

Description of the Prior Art

20

30

35

10

15

Angiogenesis is a biological process of generating new blood vessels into a tissue or organ. Under normal physiological conditions, humans or animals undergo angiogenesis only in very specific restricted situations. For example, angiogenesis is normally observed in wound healing, fetal and embryonal development and formation of the corpus luteum, endometrium and placenta. been reported that new vessel growth tightly controlled by many angiogenic regulators (see: Folkman, J., Nature Med., 1: 27-31, 1995a), and the switch of the angiogenesis phenotype depends on the net up-regulation of angiogenic stimulators and down-regulation of angiogenic suppressors.

An imbalance of the angiogenic process has been shown to contribute to pathological disorders such as diabetic retinopathy, rheumatoid arthritis and psoriasis (see: Folkman, J., Nature Med., 1: 27-31, 1995a).

15

20

25

30

Especially, both primary and metastatic tumors need to angiogenic vessels for their growth Folkman, J., New Engl. J. Med., 285:1182-1186, 1971; Folkman, J., J. Biol. Chem., 267:10931-10934, 1992). angiogenic activity could be repressed eliminated, then the tumor, although present, would not There are many reports suggesting that inhibiting tumor angiogenesis should provide a practical approach to long term control of the disease. Blocking positive regulators of angiogenesis or utilizing regulators to suppress angiogenesis results in a delay or regression of experimental tumors. If the angiogenic activity could be repressed or eliminated, then the tumor, although present, would not grow. Moreover, in the disease state, prevention of angiogenesis could avert the damage caused by the invasion of the new microvascular system effectively. Therefore, therapies directed at control of the angiogenic process could lead to the abrogation or mitigation of these diseases.

Therefore, what is needed is a novel angiogenesis inhibitor which can inhibit the unwanted growth of blood vessels, especially into tumors. An anticancer agent comprising the angiogenesis inhibitor should be able to overcome the activity of endogenous growth factors in premetastatic tumors and prevent the formation of the capillaries in the tumors thereby inhibiting the growth of the tumors. The anticancer agent should also be able to modulate the formation of capillaries in other angiogenic processes, such as wound healing reproduction. Finally, the anticancer agent and method for inhibiting angiogenesis should preferably be nontoxic and produce few side-effects.

Until now, at least 10 endogenous angiogenic inhibitors have been identified in the art (<u>see</u>: O'Reilly, M. S. et al., Cell, 88: 277-285, 1997). One such molecule is angiostatin, which consists of the plasminogen kringle I through IV(<u>see</u>: O'Reilly, M. S.

et al., Cell, 79:315-328, 1994). When applied systemically, angiostatin powerfully inhibits both primary tumor growth and metastasis without toxicity, and angiogenesis induced by bFGF as well (see: O'Reilly, M. S. et al., Nature Med., 2:689-692, 1996). These antitumor effects were accompanied by a marked reduction of microvessel density within the tumor mass, indicating that suppression of angiogenesis was associated with the inhibition of tumor growth.

Kringles are protein structural domains composed of 10 approximately 80 amino acids and three intramolecular disulfide bonds. Kringle structures are found in many proteins such as prothrombin (see: Walz, D. A. et al., Proc. Natl. Acad. Sci., U.S.A., 74:1969-1973, plasminogen(see: Ponting, C. P., Blood Coagul. & Fibrinolysis, 3:605-614, 1992), urokinase(see: Pennica, D. et al., Nature, 301:579-582 1983), hepatocyte growth factor(see: Lukker, N. A. et al., Protein Eng., 7:895-903, 1994), and apolipoprotein("apo")(a)(see: McLean, J. 20 W. et al., Nature, 330:132-137, 1987). These domains appear to be independent folding units, but their functional role is not yet known. The previous reports represent that the kringle structure can act inhibitors of endothelial cell migration and 25 proliferation during angiogenesis. Specifically, prothrombin's kringle 2 and plasminogen's kringle 1-4, and 5 have been shown to be anti-angiogenic (see: Ji, W. R. et al., FASEB J., 15:1731-1738, 1998a; Ji, W. R. et al., Biochem. Biophys. Res. Commun., 247:414-419, 1998b; 30 Cao, Y. et al., J. Biol. Chem., 271:29461-29467, 1996; Cao, Y. et al., J. Biol. Chem., 272:22924-22928, 1997; Barendsz-Janson, A. F., J. Vasc. Res., 35:109-114, 1998; Lee, T. H. et al., J. Biol. Chem., 273:28805-28812, 1998).

Apolipoprotein(a), one of the proteins having kringle structures, is a candidate for a novel angiogenesis inhibitor. Apo(a) is covalently attached to

4

apoB-100, the main protein component of low density lipoprotein(LDL) to form lipoprotein(a) (see: Fless, G. M., J. Biol. Chem., 261: 8712-8718, 1986). Elevated plasma concentration of Lp(a) represents major independent risk factor for artherosclerosis(see: Armstrong, V. W. et al., Artherosclerosis, 62:249-257, 1986; Assmann, G., Am. J. Cardiol., 77:1179-1184, 1996; Bostom, A. G. et al., JAMA, 276:544-548, 1996). Although several pathogenic activities have been reported, the physiological role of apo(a) has not yet established (see: Lawn, R. M. et al., J. Biol. Chem., 271:31367-31371, 1996; Scanu, A. M. and Fless, G. M., J. Clin. Invest., 85:1709-1715, 1990; Utermann, G., Science, 246:0904-910, 1989).

Apo(a) contains two types of kringle domains and an 15 inactive protease-like domains: the first 37 kringle domains are ~75% identical to plasminogen kringle IV, and the last kringle domain is 90% identical plasminogen kringle V. Interestingly, the kringle IV-20 like domain is present in 15-40 copies in different human alleles of the apo(a) gene. In this regard, it is feasible to develop an inhibitor of tumor angiogenesis and growth employing the Apo(a) kringle structures.

25 SUMMARY OF THE INVENTION

In accordance with the present invention, the inventors have cloned and expressed the human apo(a) kringles containing IV36, IV37 and V38 as a recombinant protein LK68, and discovered that: the LK68 protein and its single kringles, LK6, LK7 and LK8, have an ability to overcome the angiogenic activity of endogenous growth factors such as bFGF in vitro; and they may be used as active ingredients of anticancer agents.

35

30

The first object of the invention is, therefore, to provide a novel LK68 protein consisting of human apo(a)

kringle domains IV36, IV37 and V38, and cDNA encoding the LK68 protein.

The second object of the invention is to provide a novel recombinant vector containing the cDNA encoding human apo(a) kringle domains IV36, IV37 and V38.

The third object of the invention is to provide an anticancer agent which comprises the LK68 protein or its single kringles, LK6, LK7 and LK8, as an active ingredient.

The fourth object of the invention is to provide a method for treating angiogenesis-mediated disease by employing the LK68 protein.

BRIEF DESCRIPTION OF THE DRAWINGS

15

The above and the other objects and features of the present invention will become apparent from the following description given in conjunction with the accompanying drawings, in which:

20

Figure 1 is a photograph of a SDS-polyacrylamide gel electrophoresis for analysis of recombinant LK68 protein expressed in E. coli.

25

35

- Figure 2 is a photograph showing the inhibition of angiogenesis by LK68 on the chick chorioallantoic membrane (CAM).
- Figure 3(A) is a graph showing inhibition of vessel growth in the CAM as a function of LK68.
 - Figure 3 (B) is a graph showing inhibition of vessel growth in the CAM as a function of single kringles, LK6, LK7, LK8, and a control.

	Figure 4(A) is a graph showing inhibition of BCE cell proliferation by recombinant LK68 and angiostatin.
	Figure 4(B) is a graph showing inhibition of BCE cell proliferation by recombinant LK6, LK7 and LK8.
10	Figure 4(C) is a graph showing inhibition of HUVEC cell proliferation by recombinant LK68 and LK8.
15	Figure 5(A) is a graph showing BrdU labeling index of LLC cells in the presence of recombinant LK68 and LK8.
20	Figure 5(B) is a graph showing BrdU labelling index of Y1 cells in the presence of recombinant LK68 and LK8.
25	Figure 5(C) is a graph showing BrdU labelling index of TIB74 cells in the presence of recombinant LK68 and LK8.
	Figure 5(D) is a graph showing BrdU labelling index of CHO cells in the presence of recombinant LK68 and LK8.
30	Figure 5(E) is a graph showing BrdU labelling index of MSF cells in the presence of recombinant LK68 and LK8.
35	Figure 5(F) is a graph showing BrdU labelling index of NIH3T3 cells in the presence of recombinant LK68 and LK8.

- Figure 6(A) is a graph showing inhibition of HUVEC cell migration by recombinant LK68, LK8 and PK5.
- Figure 6(B) is a graph showing inhibition of HUVEC cell migration by recombinant LK68, LK6, LK7 and LK8.
- Figure 7 is a graph showing inhibition of BCE cell migration by angiostatin, recombinant LK68, LK6, LK7, and LK8 and combination of single kringles.
- Figure 8 shows the effect of administration of

 LK68 to mice having implanted

 Lewis lung carcinoma cells on

 total volume as a function of

 time.
- Figures 9(A) to 9(C) are photographs showing histological analyses of Lewis lung carcinoma cells by hematoxylin and eosin (H/E) staining.

Figure 10 shows the effect of administration of LK68 to nude mice having implanted human lung carcinoma A549 cells on total volume as a function of time.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a novel protein LK68, which can be cloned and expressed as recombinant protein from the human apolipoprotein("apo")(a) kringles. The LK68 protein consists of amino acid sequences of human

20

25

30

35

apolipoprotein(a) kringle domains IV36(amino acid 8 to 80), IV37(amino acid 122 to 194) and V38(amino acid 226 to 300) in a serial manner(see: SEQ ID NO: 2). The first two kringle domains of LK68 (i.e., IV36 and IV37) are homologous to human plasminogen kringle IV, and the third kringle domain V38 is homologous to human plasminogen kringle V. The present invention also provides a cDNA encoding the LK68 protein (see: SEQ ID NO: 1) and recombinant vectors which comprises the said cDNA and expression vectors such as pET vector series.

In describing the kringle domains of the invention, human apolipoprotein(a) kringles IV36, IV37 and V38 are abbreviated as KIV36, KIV37 and KV38, respectively; LK68 is employed to mean the recombinant protein which comprises the said three kringle domains; and, LK6, LK7 and LK8 are employed to mean the recombinant proteins of KIV36, KIV37 and KV38, respectively.

Because apolipoprotein(a) contains plasminogen-type kringle it was domains, assumed that apolipoprotein(a) could possibly have an anti-angiogenic activity. There is an experimental evidence suggesting apolipoprotein(a) may contain biological activity as an inhibitor of tumor angiogenesis and growth (see: Trieu, V. N. and Uckun, F. M., Biochem. Biophys. Res. Commun., 257:714, 1999). It has been reported that LL/2 (Lewis Lung Carcinoma) tumor growth is delayed in apo(a) transgenic mice and the microvessel density of LL/2 tumors from apo(a) transgenic mice is lower than that from wild-type mice as control.

Under the circumstance, the present inventors assumed that LK68 protein, its single kringles or their functional equivalents may have an anti-angiogenic activity. To verify said anti-angiogenic activity, it was investigated whether recombinant LK68 and its single kringles (i.e., LK6, LK7 and LK8) are potent anti-angiogenic factors in vitro and in vivo as well. As a result, LK68, LK6, LK7 and LK8 exhibit inhibitory

20

25

activities on the cultured endothelial cell proliferation as well as on the endothelial migration. LK68 and its single kringles also inhibit the normal development of capillaries in the chick embryo chorioallantoic membrane (CAM). It was also shown that systemic administration of LK68 inhibited the primary tumor growth, which is correlated with a suppression of tumor-induced angiogenesis. Since each of the single kringle proteins, LK6, LK7 LK8 and showed angiogenic activity, it is expected that they also inhibit the primary tumor growth or metastasis.

Accordingly, LK68 protein, its single kringles or their functional equivalents may be applied for the development of a potent anti-cancer agent, which is highly effective for angiogenesis-mediated diseases covering reumatoid arthritis, psoriasis, or ocular angiogenic diseases, etc.

Also, LK68 protein, its single kringles or their functional equivalents may be used in combination with other compositions and procedures for the treatment of diseases. For example, tumor may be conventionally with surgery, radiation or chemotherapy combined with LK68, its single kringles, or their functional equivalent, and then LK68, its single kringles, or their functional equivalent may subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize and inhibit the growth of any residual primary tumor.

The present invention is further illustrated in the following examples, which should not be taken to limit the scope of the invention.

Example 1: Cloning and Expression of Recombinant LK68

35

In order to verify the anti-angiogenic activity of human apo(a) kringle, the inventors cloned and expressed

25

30

35

the last three kringles containing IV36, IV37 and V38 as a recombinant protein LK68. A DNA fragment of apo(a) spanning nucleotides 12,052 to 12,975 (see: McLean J. W. et al., Nature, 330:132, 1987) was PCR-amplified from human liver cDNA and the resulting 924-bp NdeI-BamHI fragment was ligated into E.coli expression vector pET11a(Novagen, USA). The oligonucleotide primers A(SEQ ID NO: 9) and F(SEQ ID NO: 14) (see: Table 1) were used for PCR amplification under the standard PCR protocol. 10 This clone was named "pET11a/LK68", which encodes 308 amino acids including human apo(a) kringle domains, IV36, IV37 and V38(<u>see</u>: SEQ NO ID: 2). The first two kringle domains of this clone, IV36 and IV37, are homologous to human plasminogen kringle IV, and the third kringle domain V38 is homologous to human plasminogen kringle V.

The nucleotide sequences of this clone confirmed in both directions. When the nucleotide sequence of this clone was compared to the same region of the human apo(a) (see: McLean J. W. et al., Nature, 330:132, 1987), the nucleotide sequences are identical with the exception of a single base change at nucleotide 554. Our clone contains a cytosine at this position as compared to a thymidine in the sequence reported by McLean et al. (see: McLean J. W. et al., Nature, 330:132, 1987), causing an amino acid change to Thr from Met. This substitution has also been reported by other groups (see: Van der-Hoek, Y. Y. et al., Hum. Mol. Genet., 2:361-366, 1993; LoGrasso, P. V. et al., J. Biol. Chem.; 269:21820-21827, 1994) and appears to be the predominant allele for apo(a).

E.~coli~ BL21(DE3) was transformed with an expression plasmid pET11a/LK68 and recombinant LK68 protein was expressed under the following conditions. One liter of Luria-Bertani broth containing ampicillin was inoculated with 10ml of an overnight culture of E.~coli~ BL21(DE3) harboring the pET11a/LK68 plasmid and incubated with shaking at 37°C. When the OD600 of the

15

20

25

30

35

culture reached 0.4 - 0.6, isopropylthio- β -Dgalactoside(IPTG) was added at a final concentration of 1mM. Cells were grown an additional 4h after induction. Cells were harvested by centrifugation at 8000xg for These cell pellets were sonicated and the 30min at 4° C. over-expressed proteins were analyzed by SDS-PAGE(see: Figure 1). In Figure 1, Mr represents a molecular weight marker (Boehringer Mannheim, Germany); lane 1. expression of recombinant LK68 protein without IPTG induction; and, lane 2, the expression of recombinant LK68 protein with IPTG induction, respectively. Recombinant LK68 protein having a molecular weight of 37kDa was well expressed in E. coli, accumulating to about 20-30% of the total protein, as evidenced by image analysis of the scanned gel. The transformant thus prepared was designated as 'Escherichia coli BL21/LK6-8', and deposited with the Korean Collection for Cultures, #52 Oun-dong, Yusong-ku, Taejon 305 - 333, Republic of Korea, an international depository authority as accession No. KCTC0633BP on Jun. 9, 1999.

Each of single kringle domains, IV36, IV37 and V38, was cloned separately into an expression vector pET15b as described above. The oligonucleotide primers used for cloning are listed in Table 1: that is, A(SEQ ID NO: 9) and D(SEQ ID NO: 12) for KIV36 cloning; B(SEO ID NO: 10) and E(SEQ ID NO: 13) for KIV37 cloning; and, C(SEQ ID and F(SEQ ID NO: 14) for KV38 cloning, respectively. These three couples of oiligonucleotide primers were used for PCR amplification under the standard PCR protocol and the resulting clones were named "pET15b/LK6", "pET15b/LK7" and "pET15b/LK8", each of which includes the single human apo(a) kringle domains of IV36, IV37 and V38, respectively. E. coli BL21(DE3) competent cells were transformed with each of the expression plasmid, pET15b/LK6, pET15b/LK7 and pET15b/LK8. The transformant with plasmid pET15b/LK6

thus prepared was designated as 'Escherichia coli BL21(DE3)/LK6', and deposited with the Korean Collection for Type Cultures, #52 Oun-dong, Yusong-ku, Taejon 305-333, Republic of Korea, an international depository authority as accession No. KCTC0655BP on Sept. 3, 1999. The transformant with plasmid pET15b/LK7 thus prepared was designated as 'Escherichia coli BL21(DE3)/LK7', and deposited with the Korean Collection for Type Cultures the same address as above, an international depository authority as accession No. KCTC0656BP on Sept. 3, 1999. The transformant with plasmid pET15b/LK8 thus prepared was designated as 'Escherichia coli BL21/LK8', and deposited with the Korean Collection for Cultures on the same address as above, an international depository authority as accession No. KCTC0634BP on Jun. 9, 1999.

LK6, Recombinant LK7 and LK8 proteins expressed under the same conditions as fusion proteins containing N-terminal His-tag. Each of the overexpressed recombinant LK6, LK7 and LK8 protein was purified using pET His-tag system under the manufacturer's recommended condition.

Table 1. Oligonucleotide primers used for PCR cloning

25

10

15

20

Nucleotide Sequences*	Description	Location**	SEQ
			ID NO.
A. TCCATATGAAAAGCCCTGTGGTCCAGGAT	K36-5'	12052-12072	9
B. CAGTCCATATGGTCCGCCAGTGCTACCATGGCA	K37-5'	12406-12427	10
C. GGAATTCCATATGGAACAGGACTGCATGTTT	к38-5'	12718-12735	11
D. CGGGATCCTTAACCTGATTCTGTTTC	к36-3'	12310-12323	12
E. CGGGATCCTTAGACCACAGTCCCTTC	к37-3'	12658-12671	13
F. CGGGATCCTTAAGAGGATGCACA	K38-3'	12964-12975	14

^{*} Restriction sites, NdeI and BamHI are added for the cloning

13

conveniences (underlined).

** <u>See</u>: McLean *et al.*, Nature, 330:132, 1987, for nucleotide sequence(accession number is X06290).

5 Example 2: Purification of the Recombinant LK68

In order to produce the recombinant LK68, high cell-density fermentation was performed in a 5L Bioflow III bioreactor (New Brunswick Scientifics, Edison, USA) in the following medium: 4% (w/v) yeast extract, 4% (w/v) glycerol, 1% (w/v) dibasic sodium phosphate, 0.2% (w/v) monobasic potassium phosphate and $50\,\mu$ g/ml ampicillin. When the cells reached an absorbance of 100 at 600 nm, protein expression was induced with 1mM IPTG and then DO-stat fed-batch was carried out for 9h with feed media(29% (w/v) yeast extract, 39% (w/v) glycerol and 0.5% (w/v) magnesium sulfate. Cells were harvested by centrifugation at 8000xg for 30 min. Each fermentation process yielded about 80g of cell/L(wet weight).

20 To assess if LK68 was expressed in the soluble fraction or the insoluble cellular fraction of E.coli cells, the inventors analyzed the LK68 expression in these fractions. This analysis showed that LK68 was located in the insoluble cellular fraction. Thus, it was 25 necessary to denature, refold and reoxidize disulfied bonds of LK68. By using the deoxycholate and detergents, the insoluble LK68 protein purified as inclusion bodies to the extent of >95% purity. Then, the inclusion bodies were solublized with 7M urea and folded into native conformation using a 30 rapid dilution and an equilibrium dialysis scheme. the folding buffer, purified inclusion bodies were easily refolded without detectable protein aggregation. After the dialysis, the protein was purified by lysine-Sepharose 4B affinity chromatography. The protein bound 35 to lysine-Sepharose was specifically eluted by arepsilon-ACA(arepsilonamino-n-caproic acid). This suggested that the lysine-

binding site located in the KIV37 kringle of the refolded protein was fully functional. Affinity elution of LK68 with 0.1M ε -ACA yielded about 3mg of protein/g of cells(wet weight). Chromatography with polymyxin-B beads (Sigma Chemical Co., USA) was subsequently performed to eliminate any endotoxin, and residual endotoxin activity was determined with the Limulus amebocyte lysate assay kit (Biowhittaker Inc., USA). The purified protein was analyzed by SDS-PAGE and was stored at -20℃ until needed. The calculated pI value of LK68 protein is 6.13. The N-terminal amino acid sequence of the purified LK68 was confirmed by amino acid sequencing.

Example 3: Chick Chorioallantoic Membrane Assay

15

20

25

10

In order to determine whether LK68 is angiogenic in vivo, the inventors tested its ability to inhibit the development of capillaries the chorioallantoic membrane ("CAM") (see: Lee, T. H. et al., J. Biol. Chem., 273:28805-28812, 1998). Fertilized three-day-old eggs were incubated at 37° C, and a window was made after the extraction of ovalbumin. After two days of incubation, a Thermanox coverslip (Nunc Inc., USA) containing recombinant LK68 protein was applied to the CAM of individual embryos. After 48h, 20% fat emulsion was injected into the chorioallantois of the embryos, and the vessel formation around the Thermanox was examined(see: Figure 2). In Figure 2, the left photograph shows the normal development of capillaries in the CAM; and, the right shows the inhibition of angiogenesis by LK68 on CAM, respectively.

When LK68 at the dose range of 3 - $5\mu g$ was applied on the CAM, more than 60 % among the 100 eggs tested showed avascular zone around the sample applied, indicating that the growth of capillaries was inhibited. With the recombinant proteins of each kringle domain, e.g. LK6, LK7 or LK8, 60 - 70% of the eggs tested showed

25

30

inhibitory effects at the dose range of $l\mu g/CAM(\underline{see}: Figures 3(A))$ and 3(B). This in vivo study showed that apo(a) kringle domains have anti-angiogenic activity and LK68 as well as single kringle proteins is a potent inhibitor of angiogenesis. There was no evidence of toxicity in any of the chick embryos tested.

Example 4: Inhibition of Endothelial Cell Proliferation

Recombinant LK68, LK6, LK7 and LK8 proteins were assayed for their inhibitory activity on proliferation of bovine capillary endothelial (BCE) cells stimulated by bFGF under the following conditions. BCE cells were grown in DMEM containing 10% bovine calf serum (BCS) and 3 ng/ml bFGF(Upstate Biotechnology, USA). Approximately 3,000 cells were added to each well of 96-well tissue culture plate and incubated at 37 °C in 5% CO₂ atmosphere. After incubation for 18 h, the medium was replaced with DMEM containing 0.5% BCS, and the test samples were added to each well. After 30 min incubation, bFGF was added to a final concentration of lng/ml. The cell count was determined by [³H]thymidine incorporation method. The experiments were performed in triplicate.

As can be seen in Figure 4, it was determined that LK68, LK6, LK7 and LK8 specifically inhibited BCE cell proliferation in a dose-dependent manner. When the angiostatin was applied as a positive control, all the Apo(a) kringle proteins tested appeared to be more effective under the conditions used in this experiment. The concentration of half-maximal inhibition (ED $_{50}$) for LK68 is determined about 200 - 250nM, about 140 - 170nM for LK6, about 10 - 20nM for LK7, and about 10 - 20nM for LK8 (see: Figures 4(A) and 4(B)).

Recombinant LK68 and LK8 proteins were assayed for their inhibitory activity on proliferation of human umbilical vein endothelial (HUVEC) cells stimulated by bFGF under the following conditions. HUVECs(American

35

Type Culture Collection, USA) were grown in F12K medium containing 10% heat-inactivated fetal bovine serum("FBS")(Hyclone, USA), $30 \,\mu\,\mathrm{g/ml}$ endothelial cell growth supplement(ECGS)(Sigma Chemical Co., USA), $100 \,\mu\,\mathrm{g/ml}$ heparin(Sigma Chemical Co., USA). The cells were plated at a density of 2000/well in 96-well tissue culture plate. The cells were incubated at 37° C, 5% CO_2 , for 18hr, washed once with serum-free medium, and F12 medium containing 0.5% FBS was added. The cells were treated with various concentrations of 10 samples and for incubated 30min. Then, ECGS, heparin and bFGF(Upstate Biotechnology, USA) were added into the cells with the final concentrations of $30\,\mu\,\mathrm{g/ml}$, $100\,\mu$ g/ml and 5ng/ml, respectively. After 48hr of incubation, cell counts were determined with the Cell Proliferation ELISA using 5-bromo-2'- deoxyuridine (BrdU) (Boehringer Mannheim, USA). The experiments were performed in triplicate.

As can be seen in Figure 4(C), it was determined that LK68 as well as LK8 specifically inhibited HUVEC cell proliferation in a dose-dependent manner.

In the presence of LK68 or single kringle proteins such as LK6, LK7 and LK8, the morphology of BCE or HUVEC cells appeared similar to those of untreated cells. addition, cell proliferation can be rescued with bFGF stimulation after removal of LK68. These indicate that LK68 as well as single kringle proteins cytotoxic to capillary endothelial Furthermore, the inhibitory activity would appear to be specific for endothelial cells, e.g., BCE and HUVEC cells. Additionally, LK68 as well as LK8 failed to show inhibition of proliferation of non-endothelial types, such as CHO cells, mouse skin fibroblast NIH3T3 cells, mouse Lewis lung carcinoma cells, mouse adrenal tumor Y1 cells and mouse embryonic liver/SV40 transformed cell line TIB74(see: Figures 5(A) to 5(F)). Figures 5(A) to 5(C) represent the sensitivity of

various tumor cells such as LLC, Y1, and TIB 74, and Figures 5(D) to 5(F) represent the sensitivity of various normal cell lines such as CHO, MSF, and NIH3T3, respectively.

5

10

15

20

25

35

Example 5: Inhibition of Endothelial Cell Migration

Cell migration assay was performed in Transwells with 8-mm pores (Costar, USA). Briefly, the wells were coated with fibronectin $(25 \mu g/ml)$ (Sigma Chemical Co., USA) overnight and HUVECs were plated at a density of 2000/well in 100 μ l Dulbecco's modified Eagle's medium containing 0.4% fetal calf serum(FCS) in the upper chamber. $500\,\mu\,\mathrm{l}$ of DMEM containing 0.4% FCS was added to the lower chamber and incubated at 37° C for 1 hr. test samples of $1\mu M$ concentration were added to the upper chamber and 25 ng/ml of bFGF was added to the lower chamber. After 5 hr incubation, cells that crossed the fibronectin-plated membrane were quantified after wiping off the cells in the upper chamber with a cotton The cells across the membrane were stained with Diff-Quik stain set according to the manufacturer's instruction (Dade Behring Inc., USA) and were counted at 100x magnification. The experiments were performed in duplicate.

Basic FGF(25ng/ml) was used to stimulate the migration of HUVEC cells. With the dose of $1\,\mu\text{M}$, LK68 as well as single kringle proteins such as LK6, LK7 and LK8 completely inhibited the bFGF-induced HUVEC cell migration to the level of uninduced control(see: Figures 6(A) and 6(B)). In Figure 6, (-)CON represent uninduced control, and (+)CON represent bFGF-induced positive control.

Migration assay using BCE cells was performed as described above. Two different concentrations of LK68 or single kringle proteins applied and all the Apo(a) kringle proteins tested showed inhibitory effects on BCE

cell migration. In addition, LK68 and its single kringle proteins were more effective on the inhibition of BCE cell migration than angiostatin(AS)(see: Figure 7).

5 Example 6: Suppression of Primary Tumor Growth

Example 6-1: Lewis Lung Carcinoma

Male 6 to 8-week-old C57BL6/J mice were implanted with Lewis lung carcinomas. The subcutaneous dorsa of mice in the proximal midline were injected with 1×10^6 cells in 0.1ml of saline. When the tumors reached about 5mm in diameter, tumor-bearing mice received LK68(100 mg/kg body weight) as a suspension in PBS injected 15 subcutaneously at a site distant from the tumor. control group of mice had only a sham procedure and was treated with PBS only. Tumor size was measured every day during the treatment; and, volumes were determined using the formula width 2 x length x 0.52 and the ratio of treated to control tumor volume(T/C) was determined for the last time point. Treatments were continued for 8 days, at which point all mice were sacrificed and the tumors were removed(see: Figure 8). As can be seen in Figure 8, it was clearly determined that the growth of LLC primary tumors was potently suppressed by systemic LK68 therapy; LK68 at a dose of 100mg/kg caused significant regression of tumor burden only with 7 day treatment.

Histological analyses were also carried out to compare tumors from treated and control mice in terms of vessel density and hemorrhage formation, and morphological appearance (see: Figures 9(A) to 9(C)). In Figures 9(A) to 9(C), 9(A) shows PBS-treated control, 9(B) LLC tumors of 10mg/kg body weight LK68-treated, and 9(C) LLC tumor of 100mg/kg body weight LK68-treated, respectively. Obvious histological differences were observed in LK68-treated tumors by hemotoxylin and

30

eosin(H/E) staining: that is tumor cells were not intact and morphologically not viable; and, zonal necrosis was examined around the tumors. Also, vessel density within LK68-treated tumors was reduced. There was no evidence of inflammation or bleeding in any of the mice treated with the recombinant LK68.

Example 6-2: Human Lung Carcinoma

Four-week-old outbred female nu/nu nude mice used 10 in this experiment were housed in a sterile environment. Cages, bedding, food and water were all autoclaved. mice were maintained on a 12-hr light/ 12-hr dark cycle. Human lung cancer cells (A549 purchased from Korean Cell Line Bank) were maintained in RPMI 1640 15 supplemented with 10% heat-inactivated **FBS** antibiotics. Approximately 2 x 107 cells of A549 human lung carcinoma were subcutaneously injected into nude mice into the proximal midline of the dorsa. When tumors were palpable at day 7 after tumor implantation, the mice were treated with LK68 at the dose of 100mg/kg body The control group was treated with PBS only. weight. The treatment was continued for 17 days. The tumor size was measured every other day.

The tumor growth was regressed by the LK68 treatment: that is, LK68-treated A549 tumors were approximately 57.5% smaller than tumors in control animals(see: Figure 10). There was no evidence of any toxicity in any of the treated mice. Continued therapy maintained the tumors in a state of dormancy for as long as it was administered. These data strongly suggest that the anti-angiogenic effect of LK68 can be used to target a wide variety of primary malignancies.

As clearly illustrated and demonstrated as above, the present invention provides a novel angiogenesis inhibitor, LK68 whose amino acid sequence is identical

with the human apo(a) kringle domains IV36, IV37 and V38, a DNA sequence encoding the LK68, a recombinant expression vector comprising the DNA, a recombinant microorganism transformed with the recombinant expression vector, use of the LK68 as an anticancer agent, and a method for treating angiogenesis-mediated disease.

BUDAPERT TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROCOLONISMS FOR THE PURPOSE OF FATENT PROCEDURE

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT

issued pursuant to Rule 7.1

TO: Mogern Biotechnology Research Institute
#241, Pojungeri, Koosing-myrm, Yongin-si, Kyongsi-do 449-910,
Republic of Kores

I, IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR:

Escherichia cali BL21(DXI)/LEE Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY:

KCTC 0855BP

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by

[x] a scientific description

[] a proposed taxonomic designation (Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Sep 03 1999.

IV RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this international Depositary Authority on and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name Korean Collection for Type Cultures

Address: Korea Research Institute of Bioscience and Biotechnology (KRIBB)

#62, Oun=dong, Yusong=ku, Taajon 306-333,

Republic of Kores

Signature(s) of person(s) having the power to represent the International Depositary Authority of authorized official(a):

BAE, Kyung Sook, Director Date: Sep 08 1999 BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROGRAMMAS FOR THE PURPOSE OF PATENT PROCEDURE

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT

issued pursuant to Rule 7.1

TO: Mogern Biotechnology Research Institute #341, Pojunu-ri, Koosung-myun, Yangin-si, Kyanggi-da 449-910, Republic of Kores

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR:

Escherichia coli HL21(DS3)/LK7 Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY:

KCTC 0856BP

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

[x] a scientific description

[) a proposed taxonomic designation

(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Sep 03 1929.

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary

Authority on and a request to convert the original deposit to a deposit

under the Budapest Treaty was received by it on

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Korean Collection for Type Cultures

Address: Korea Research Institute of Buscience and Blotechnology (KRIBB)

#52, Oun-dong, Yusong-ku,

Tanjon 305-333, Republic of Korea Signature(s) of person(s) having the power to represent the international Depositary Authority of authorized official(s):

BAE, Kyung Sook, Director Date: Sep 08 1999

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSE OF PATENT PROCEDURE

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT

issued oursuant to Rule 7.1

70 Mogam Biotech, Research Institute #341, Philung-ri, Koosung-myun, Yongin-si, Kyonggi-do 449-910, Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR

Escherichia coli BL21/LK8

Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY:

KCTC 0634BP

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

{ x } a scientific description

[] a proposed taxonomic designation (Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above, which was received by it on **Jun 09 1999**.

N RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary Authority on and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Korean Collection for Type Cultures

Address Korea Research Institute of Bioscience and Biotechnology

(KRIBB)

#52, Oun-dong, Yusong-ku,

Taejon 305-333, Republic of Korea Signature(s) of person(s) having the power to represent the International Depositary Authority of authorized official(s):

BAE, Kyung Sook, Director Date' Jun 16 1999

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSE OF PATENT PROCEDURE

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT

issued pursuant to Rule 7.1

TO Mogam Biotech. Research Institute #341, Popung-ri, Koosung-myun, Yongin-si, Kyonggi-do 449-910, Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR:

Escherichia coli BL21/LK6-8

Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY

KCTC 0633BP

II SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by [\mathbf{x}] a scientific description

[] a proposed taxonomic designation (Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above, which was received by it on **Jun 09 1999**.

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary Authority on and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name. Korean Collection for Type Cultures

Address: Korea Research Institute of Bioscience and Biotechnology

(KRIBB)

#52, Oun-dong, Yusong-ku.

Taejon 305-333, Republic of Korea Signature(s) of person(s) having the power to represent the International Depositary Authority of authorized official(s):

BAE, Kyung Sook, Director Date: Jun 16 1999

WHAT IS CLAIMED IS:

- LK6 protein (SEQ ID NO: 4) consisting of amino acid sequences of human apolipoprotein(a) kringle domains IV36.
 - 2. LK7 protein (SEQ ID NO: 6) consisting of amino acid sequences of human apolipoprotein(a) kringle domains IV37.

10

- 3. LK8 protein (SEQ ID NO: 8) consisting of amino acid sequences of human apolipoprotein(a) kringle domains V38.
- 4. LK68 protein (SEQ ID NO: 2) consisting of amino acid sequences of human apolipoprotein(a) kringle domains IV36, IV37 and V38 in a serial manner
- 5. A cDNA sequence (SEQ ID NO: 3) which codes for the LK6 protein of claim 1.
 - 6. A cDNA sequence(SEQ ID NO: 5) which codes for the LK7 protein of claim 2.
- 7. A cDNA sequence(SEQ ID NO: 7) which codes for the LK8 protein of claim 3.
 - 8. A cDNA sequence(SEQ ID NO: 1) which codes for the LK68 protein of claim 4.

- 9. A recombinant expression vector pET15b/LK6 comprising the cDNA of claim 5 which expresses the LK6 protein of claim 1.
- 10. A recombinant expression vector pET15b/LK7 comprising the cDNA of claim 6 which expresses the LK7 protein of claim 2.

11. A recombinant expression vector pET15b/LK8 comprising the cDNA of claim 7 which expresses the LK8 protein of claim 3.

5

- 12. A recombinant expression vector pET11a/LK68 comprising the cDNA of claim 8 which expresses the LK68 protein of claim 4.
- 13. Escherichia coli BL21(DE3)/LK6(KCTC0655BP) transformed with the recombinant expression vector pET15b/LK6 of claim 9.
- 14. Escherichia coli BL21(DE3)/LK7(KCTC0656BP)

 15 transformed with the recombinant expression vector pET15b/LK7 of claim 10.
- 15. Escherichia coli BL21/LK8(KCTC0634BP) transformed with the recombinant expression vector pET15b/LK8 of claim 11.
 - 16. Escherichia coli BL21/LK6-8(KCTC0633BP) transformed with the recombinant expression vector pET11a/LK68 of claim 12.

25

17. An anticancer agent which comprises an active ingredient of LK68 protein, its single kringles, or their functional equivalents and pharmaceutically acceptable carrier.

30

18. A method for treating angiogenesis-mediated disease which comprises administering therapeutically effective amount of LK 68 protein, its single kringles, or their functional equivalents to a human or animal.

35

19. The method for treating angiogenesis-mediated disease of claim 18, wherein the angiogenesis-mediated

disease is cancer, rheumatoid arthritis, psoriasis, or ocular angiogenic disease.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 22 March 2001 (22.03.2001)

PCT

(10) International Publication Number WO 01/19868 A1

(51) International Patent Classification?: C12N 15/63, 15/70, 1/21, A61K 38/17 C07K 14/47,

- (21) International Application Number: PCT/KR99/00554
- (22) International Filing Date:

15 September 1999 (15.09.1999)

(25) Filing Language:

English

(26) Publication Language:

English

- (71) Applicant (for all designated States except US): MOGAM BIOTECHNOLOGY RESEARCH INSTITUTE [KR/KR]; 341, Pojung-ri, Koosung-myon, Yongin-si, Kyonggi-do 449-910 (KR).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): CHANG, Jihoon [KR/KR]; 553-517, Jayang 3-dong, Kwangjin-ku, Seoul 143-193 (KR). KIM, Jang, Seong [KR/KR]; 902 Ikjoo Apt., Yeonmoo-dong, JangAn-ku, Suwon-si, Kyonggi-do 440-240 (KR). PARK, Eun, Jeong [KR/KR]; 322-1007, Woosung Apt., Sangrok-maeul, Jungja-dong, Pundang-ku, Sungnam-si, Kyonggi-do 464-010 (KR). YUM, Jungsun [KR/KR]; 115-502, Chunggu Apt., Hansol-maeul, Jungja-dong, Pundang-ku, Sungnam-si, Kyonggi-di

463-010 (KR). CHUNG, Soo-il [KR/KR]; 112-902, Hyundae Apt., Hyoja-chon, Seohyun-dong, Pundang-ku, Sungnam-si, Kyonggi-do 463-480 (KR).

- (74) Agent: LEE, Han-Young; Seowon Building, 8th floor, 1675-1, Seocho-dong, Seocho-gu, Seoul 137-070 (KR).
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A NOVEL ANGIOGENESIS INHIBITOR

(57) Abstract: The present invention provides a novel angiogenesis inhibitor, LK68 whose amino acid sequence is identical with the human apolipoprotein (a) kringle domains IV36, IV37 and V38, a cDNA sequence encoding the LK68, a recombinant expression vector comprising the cDNA, a recombinant microorganism transformed with the recombinant expression vector and a novel use of the LK68 as an anticancer agent and a method for treating angiogenesis-mediated disease. LK68, LK6, LK7 and LK8 exhibit inhibitory activities on the cultured endothelial cell proliferation as well as on the endothelial cell migration. LK68 and its single kringles also inhibit the normal development of capillaries in the chick embryo chorioallantoic membrane (CAM). It was also showed that systemic administration of LK68 causes the inhibition of primary tumor growth, which is correlated with a suppression of tumor-induced angiogenesis. Accordingly, LK68 protein, its single kringles or their functional equivalents may be applied for the development of a potent anti-cancer agent, which is highly effective for angiogenesis-mediated diseases covering cancer, rheumatoid arthritis, psoriasis, ocular angiogenic disease, etc.





FIG. 1

PCT/KR99/00554






FIG. 2






FIG. 3(A)




FIG. 3(B)




FIG. 4(A)




FIG. 4(B)




FIG. 4(C)

.....




FIG. 5(A)




FIG. 5(B)




FIG. 5(C)




FIG. 5(D)




FIG. 5(E)




FIG. 5(F)




FIG. 6(A)




FIG. 6(B)




FIG. 7

Suppression of LLC by LK68




FIG. 8




FIG. 9(A)




FIG. 9(B)



FIG. 9(C)

Tumor suppression by LK 6-8 (hlung carcinoma)




FIG. 10

*Express Mail Label No. EL 824098099 US

Patent 58049-00002

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare:

THAT my residence, post office address and citizenship are as stated below next to my name.

THAT I believe I am the original, first and sole (if only one name is listed below) or an original, first and joint inventor (if plural inventors are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: A NOVEL ANGIOGENESIS INHIBITOR the specification of which:

	[X] is attached hereto.
	OR	
	[] was filed on, as United States Application Number
		·
THAT	the	subject matter of the
	[] attached amendment
	OR	
	[] amendment filed on (MM/DD/YY)

was part of my or our invention and was invented before the filing date of the original application, above identified for such invention.

THAT I do not know and do not believe that this invention was ever known or used in the United States of America before my or our invention or discovery thereof, or patented or described in any printed publication in any country before my or our invention or discovery thereof, for more than one year prior to this application.

THAT the invention was not in public use or on sale in the United States of America for more than one year prior to this application.

THAT this invention has not been patented or made the subject of an inventor's certificate issued before the date of the application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months before this application.

THAT I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

THAT I acknowledge the duty to disclose information of which I am aware which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations §1.56.

THAT no application(s) for patent or inventor's certificate on this invention or discovery has been filed by me or my legal representatives or assigns in a country foreign to the United States of America more than 12 months prior hereto, unless identified here:

NONE

.

THAT I hereby claim foreign priority benefits under Title 35, United States Code §119 (a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application	Country	Foreign Filing Date	Priority Claimed?	Certified Copy Attached?		
Number(s)		(MM/DD/YY)	YES	YES		
		ł	NO	NO		
PCT/KR99/00554	PCT	September 15, 1999	X	X		

THAT I hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YY)

THAT I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s), or §365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

Number		(MM/DD/YY)	(If applicable)
U.S. Parent Application	PCT Parent Number	Parent Filing Date	Parent Patent Number

Express Mail Label No. EL 824098099 US

Patent 58049-00002

And as a named inventor, I hereby appoint the following registered practitioners to prosecute this application and to transact all business in the United States Patent and Trademark Office connected therewith and with the resulting patent, individually and collectively:

Squire, Sanders & Dempsey L.L.P. 801 S. Figueroa Street, 14th Floor Los Angeles, California 90017-5554

telephone number (213) 624-2500 (to whom all communications regarding the subject application are to be directed); and each practitioner thereof named below with Registration Numbers, and of the same address:

Joseph Hyosuk Kim, Ph. D.

Reg. No. 41,425

and further appoint as associate practitioners, with right of revocation in the primary practitioners, the following:

Q

Michael A. Lechter, Reg. No. <u>27,350</u> Marc A. Sockol, Reg. No. <u>40,823</u> William R. Bachand, Reg. No. <u>34,980</u> David B. Abel, Reg. No. <u>32,394</u> David E. Rogers, Reg. No. 38,287 Lorinda J. Howland, Reg. No. 42,671 Dave B. Koo, Reg. No. 46,839 Sung I. Oh, Reg No. 45,583

whose address is:

SQUIRE, SANDERS & DEMPSEY L.L.P 801 So. Figueroa St., 14th Fl. Los Angeles, CA 90017-5554

Telephone: (213) 624-2500

Please direct all correspondence to:

Joseph Hyosuk Kim, Ph.D. Squire, Sanders & Dempsey L.L.P. 801 S. Figueroa Street, 14th Floor Los Angeles, California 90017-5554

I hereby declare further that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Patent 58049-00002

Sant All the second of the sec	
Inventor's Signature	
Date March 7.2002	
Inventor's Name (typed) Jihoon Chang First Middle Family Name	
Citizenship Korean	
Post Office Address 553-517, Jayang 3-dong, Kwangjin-ku	
Residence (City) Seoul	
(State/Foreign Country) Korea (Zip Code) 143-193	
Inventor's Signature J. S. Kim	
Date March 7. 2002	
Inventor's Name (typed) Jang Seong Kim First Middle Family Name	
Citizenship Korean	
Post Office Address 902 Ikjoo Apt., Yeonmoo-dong, JangAn-ku	
Residence (City) Suwon-si	
(State/Foreign Country) Kyonggi-do, Korea (Zip Code) 440-240	
Inventor's Signature 184 2 76	
Date March 7 2002	
Inventor's Name (typed) 2 Eun Jeong Park First Middle Family Name	
Citizenship Korean	
Post Office Address 322-1007, Woosung Apt., Sangrok-maeul, Jungja-dong, Pundang-ku	K
Residence (City) Sungnam-si	
(State/Foreign Country) Kyonggi-do, Korea (Zip Code) 464-010	

Express Mail Label No. EL 824098099 US

Patent 58049-00002

Inventor's Signature	Jungsun	Yun		
Date March	7 2002			
Inventor's Name (typed)	First Jung-sun	Middle	Yum Family Name	
Citizenship Korean	-			
Post Office Address	115-502 Chunggu Apt.,	Hansol-maeul, Jungja-	dong, Pundang-ku	KR.
Residence (City)	Sungnam-si			
(State/Foreign Country)	Kyonggi-do, Korea	_(Zip Code) <u>463-010</u>	<u> </u>	
Inventor's Signature	Sooll	Chung		
Date <u>Vareh</u>	2. 2002	đ		
Inventor's Name (typed)_	Soo-il First	Middle	Chung Family Name	
Citizenship Korean		Middle	ranny Name	
Post Office Address	112-902 Hyundae Apt.,	Hyoja-chon, Seohyun-	dong, Pundang-ku	HRY
Residence (City)	Sungnam-si			., 9
(State/Foreign Country)	Kyonggi-do Korea	_(Zip Code) <u>463-480</u>	<u>i </u>	•

JC10 Rec'd PCT/PTO 1 5 MAR 2002/

SEQUENCE LISTING

	1102	mogam broteen	iorogy kese				
5	<120>	A NOVEL ANGIO	GENESIS INH	IBITOR			
	<160>	14					
10	<170>	KOPATIN 1.5					
	<210>	1					
	<211>	924					
	<212>	DNA					
15	<213>	Homo sapiens					
	<400>	1					
		tg tggtccagga	ttgctaccat	ggtgatggac	ggagttatcg	aggcatatcc	60
20	tccaccac	tg tcacaggaag	gacctgtcaa	tcttggtcat	ctatgatacc	acactggcat	120
	cagaggac	cc cagaaaacta	cccaaatgct	ggcctgaccg	agaactactg	caggaatcca	180
25	gattctgg	ga aacaaccctg	gtgttacaca	accgatccgt	gtgtgaggtg	ggagtactgc	240
23	aatctgac	ac aatgctcaga	aacagaatca	ggtgtcctag	agactcccac	tgttgttcca	300
	gttccaag	ca tggaggctca	ttctgaagca	gcaccaactg	agcaaacccc	tgtggtccgc	360
30	cagtgcta	cc atggcaatgg	ccagagttat	cgaggcacat	tctccaccac	tgtcacagga	420
	aggacatg	tc aatctiggic	atccatgaca	ccacaccggc	atcagaggac	cccagaaaac	480
35	tacccaaa	tg atggcctgac	aatgaactac	tgcaggaatc	cagatgccga	tacaggccct	540
35	tggtgttt	ta ccacggaccc	cagcat cagg	tgggagtact	gcaacctgac	gcgatgctca	600

	gacac	aga	ag g	gact	gtgg	gt cg	gctco	ctccs	g act	gtca	tcc	aggt	tcca	ag (ect ag	gggcc	:t	660
	ccttc	etga	ac a	agao	etgta	at g	tttg	ggaa	ggg	aaag	gat	accg	gggg	caa g	gaagg	gcaac	cc	720
5	actgt	tac	tg g	gacg	gccai	tg co	cagga	aatg	g gct	gccc	agg	agco	ccat	ag a	acaca	agcac	g	780
	ttcat	tcc	ag g	gaca	aata	aa at	t ggg	caggi	ctg	gaaa	aaa	atta	actgo	ccg	taaco	cctga	ıt	840
10	ggtga	acat	ca a	itggi	ccci	tg gi	tgct	acac	a atg	gaatc	caa	gaaa	acti	itt 1	tgact	tactg	ţt	900
	gatat	ccc	tc t	ctgt	gca	tc ci	tct								•			924
	<210>	>	2															
15	<211>		308	}														
	<212>		PRI															
	<213>				ap i er	าร												
	<400>	•	2															
20																		
	Lys S	Ser I	Pro	Val	Val	Gln	Asp	Cys	Tyr	His (Gly	Asp	Gly	Arg	Ser	Tyr		
	Lys S	Ser I	Pro	Val	Val 5	Gln	Asp	Cys	Tyr	His (Gly	Asp	Gly	Arg	Ser 15	Tyr		
	Lys S	Ser I	Pro	Val		Gln	Asp	Cys	Tyr		Gly	Asp	Gly	Arg		Tyr		
					5					10					15			
	1				5					10					15			
25	1			Ser	5				Thr	10				Gln	15			
	1	lly i	lle	Ser 20	5 Ser	Thr	Thr	Val	Thr 25	10 Gly 1	Arg	Thr	Cys	G1n 30	15 Ser	Trp		
	1 Arg G	lly i	lle	Ser 20	5 Ser	Thr	Thr	Val	Thr 25	10 Gly 1	Arg	Thr	Cys	G1n 30	15 Ser	Trp		
	1 Arg G	lly i	Ile Met	Ser 20	5 Ser	Thr	Thr	Val His	Thr 25	10 Gly 1	Arg	Thr	Cys Glu	G1n 30	15 Ser	Trp		
	1 Arg G	Gly Ser l	Ile Met 35	Ser 20 Ile	5 Ser Pro	Thr	Thr Trp	Val His 40	Thr 25 Gln	10 Gly A	Arg Thr	Thr Pro	Cys Glu 45	G1n 30 Asn	15 Ser Tyr	Trp Pro		
	1 Arg G Ser S	Gly Ser l	Ile Met 35	Ser 20 Ile	5 Ser Pro	Thr	Thr Trp Asn	Val His 40	Thr 25 Gln	10 Gly A	Arg Thr	Thr Pro	Cys Glu 45	G1n 30 Asn	15 Ser Tyr	Trp Pro		
25	1 Arg G Ser S	Gly Ser I	Ile Met 35	Ser 20 Ile	5 Ser Pro	Thr His Glu	Thr Trp Asn	Val His 40	Thr 25 Gln	10 Gly A	Arg Thr	Thr Pro Pro	Cys Glu 45	G1n 30 Asn	15 Ser Tyr	Trp Pro		
25	1 Arg G Ser S	Gly Ger ! Mla (Ile Met 35 Gly	Ser 20 Ile Leu	5 Ser Pro	Thr His Glu	Thr Trp Asn 55	Val His 40 Tyr	Thr 25 Gln Cys	10 Gly A Arg	Arg Thr Asn	Thr Pro Pro 60	Cys Glu 45 Asp	G1n 30 Asn Ser	15 Ser Tyr Gly	Trp Pro Lys		
25	Arg G Ser S Asn A	Gly Ger ! Mla (Ile Met 35 Gly	Ser 20 Ile Leu	5 Ser Pro	Thr His Glu	Thr Trp Asn 55	Val His 40 Tyr	Thr 25 Gln Cys	10 Gly A Arg	Arg Thr Asn	Thr Pro Pro 60	Cys Glu 45 Asp	G1n 30 Asn Ser	15 Ser Tyr Gly	Trp Pro Lys		
25	Arg G Ser S Asn A Gln P	Gly Ser I	Met 35 Gly	Ser 20 Ile Leu Cys	5 Ser Pro Thr	Thr His Glu Thr 70	Thr Trp Asn 55	Val His 40 Tyr	Thr 25 Gln Cys	10 Gly Arg	Arg Thr Asn Val 75	Thr Pro Pro 60 Arg	Cys Glu 45 Asp Trp	Gln 30 Asn Ser	15 Ser Tyr Gly	Trp Pro Lys Cys 80		
25	Arg G Ser S Asn A	Gly Ser I	Met 35 Gly	Ser 20 Ile Leu Cys	5 Ser Pro Thr	Thr His Glu Thr 70	Thr Trp Asn 55	Val His 40 Tyr	Thr 25 Gln Cys	10 Gly Arg	Arg Thr Asn Val 75	Thr Pro Pro 60 Arg	Cys Glu 45 Asp Trp	Gln 30 Asn Ser	15 Ser Tyr Gly	Trp Pro Lys Cys 80		

•	Thr	Val	Val	Pro 100	Val	Pro	Ser	Met	Glu 105	Ala	His	Ser	Glu	Ala 110	Ala	Pro
5	Thr	Glu	Gln 115	Thr	Pro	Val	Val	Arg 120	Gln	Cys	Tyr	His	Gly 125	Asn	Gly	Glr
	Ser	Tyr 130	Arg	Gly	Thr	Phe	Ser 135	Thr	Thr	Val	Thr	Gly 140	Arg	Thr	Cys	Glr
10	Ser 145	Trp	Ser	Ser	Met	Thr 150	Pro	His	Arg	His	Gln 155	Arg	Thr	Pro	Glu	Asr 160
15	Tyr	Pro	Asn	Asp	Gly 165	Leu	Thr	Met	Asn	Tyr 170	Cys	Arg	Asn	Pro	Asp 175	Ala
13	Asp	Thr	Gly	Pro 180	Trp	Cys	Phe	Thr	Thr 185	Asp	Pro	Ser	He	Arg 190	Trp	Glu
20	Tyr	Cys	Asn 195	Leu	Thr	Arg	Cys	Ser 200	Asp	Thr	Glu	Gly	Thr 205	Val	Val	Ala
	Pro	Pro 210	Thr	Val	He	Gln	Val 215	Pro	Ser	Leu	Gly	Pro 220	Pro	Ser	Glu	Gli
25	Asp 225	Cys	Met	Phe	Gly	Asn 230	Gly	Lys	Gly	Tyr	Arg 235	Gly	Lys	Lys	Ala	Thi 240
30	Thr	Val	Thr	Gly	Thr 245	Pro	Cys	Gln	Glu	Trp 250	Ala	Ala	Gln	Glu	Pro 255	His
30	Arg	His	Ser	Thr 260	Phe	He	Pro	Gly	Thr 265	Asn	Lys	Trp	Ala	Gly 270	Leu	Gli
35	Lys	Asn	Tyr 2 7 5	Cys	Arg	Asn	Pro	Asp 280	Gly	Asp	Ile	Asn	Gly 285	Pro	Trp	Cy
	Tyr	Thr	Met	Asn	Pro	Arg	Lys	Leu	Phe	Asp	Tyr	Cys	Asp	Ile	Pro	Lei

295 300 290 Cys Ala Ser Ser 305 5 3 <210> <211> 273 <212> DNA Homo sapiens <213> 10 <400> 3 aaaagccctg tggtccagga ttgctaccat ggtgatggac ggagttatcg aggcatatcc 60 tccaccactg tcacaggaag gacctgtcaa tcttggtcat ctatgatacc acactggcat 120 15 cagaggaccc cagaaaacta cccaaatgct ggcctgaccg agaactactg caggaatcca 180 gattctggga aacaaccctg gtgttacaca accgatccgt gtgtgaggtg ggagtactgc 240 20 273 aatctgacac aatgctcaga aacagaatca ggt <210> 4 <211> 91 25 <212> PRT <213> Homo sapiens <400> Lys Ser Pro Val Val Gln Asp Cys Tyr His Gly Asp Gly Arg Ser Tyr 30 1 Arg Gly Ile Ser Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ser Trp

25

45

Ser Ser Met Ile Pro His Trp His Gln Arg Thr Pro Glu Asn Tyr Pro 40

20

35

	Asn Ala Gly Leu Thr Glu Asn Tyr Cys Arg Asn Pro Asp Ser Gly Lys 50 55 60	
5	Gln Pro Trp Cys Tyr Thr Thr Asp Pro Cys Val Arg Trp Glu Tyr Cys 65 70 75 80	
	Asn Leu Thr Gln Cys Ser Glu Thr Glu Ser Gly 85 90	
10		
15	<210> 5 <211> 267 <212> DNA <213> Homo sapiens	
	44005	
	<400> 5 gtccgccagt gctaccatgg caatggccag agttatcgag gcacattctc caccactgtc	60
20	acaggaagga catgtcaatc ttggtcatcc atgacaccac accggcatca gaggacccca	120
	gaaaactacc caaatgatgg cctgacaatg aactactgca ggaatccaga tgccgataca	180
25	ggcccttggt gttttaccac ggaccccagc atcaggtggg agtactgcaa cctgacgcga	240
23	tgctcagaca cagaagggac tgtggtc .	267
	<210> 6	
30	<211> 89	
	<212> PRT <213> Homo sapiens	
	<400> 6	
35	Val Arg Gln Cys Tyr His Gly Asn Gly Gln Ser Tyr Arg Gly Thr Phe	
	1 5 10 15	

Ser Thr Thr Val Thr Gly Arg Thr Cys Gln Ser Trp Ser Ser Met Thr 30 25 20 Pro His Arg His Gln Arg Thr Pro Glu Asn Tyr Pro Asn Asp Gly Leu 35 40 5 Thr Met Asn Tyr Cys Arg Asn Pro Asp Ala Asp Thr Gly Pro Trp Cys **6**0 55 50 Phe Thr Thr Asp Pro Ser Ile Arg Trp Glu Tyr Cys Asn Leu Thr Arg 10 70 75 Cys Ser Asp Thr Glu Gly Thr Val Val 85 15 <210> 7 <211> 258 <212> DNA <213> Homo sapiens 20 <400> 7 60 gaacaggact gcatgtttgg gaatgggaaa ggataccggg gcaagaaggc aaccactgtt 120 25 actgggacgc catgccagga atgggctgcc caggagcccc atagacacag cacgttcatt ccagggacaa ataaatgggc aggtctggaa aaaaattact gccgtaaccc tgatggtgac 180 240 atcaatggtc cctggtgcta cacaatgaat ccaagaaaac tttttgacta ctgtgatatc 30 258 cctctctgtg catcctct <210> 8 35 <211> 86 <212> PRT <213> Homo sapiens

<400> 8

Glu Gln Asp Cys Met Phe Gly Asn Gly Lys Gly Tyr Arg Gly Lys Lys

1

5

10

15

5

Ala Thr Thr Val Thr Gly Thr Pro Cys Gln Glu Trp Ala Ala Gln Glu
20 25 30

Pro His Arg His Ser Thr Phe Ile Pro Gly Thr Asn Lys Trp Ala Gly

10 3

40

45

Leu Glu Lys Asn Tyr Cys Arg Asn Pro Asp Gly Asp IIe Asn Gly Pro
50 55 60

Trp Cys Tyr Thr Met Asn Pro Arg Lys Leu Phe Asp Tyr Cys Asp IIe
65 70 75 80

Pro Leu Cys Ala Ser Ser

85

20

<210> 9

<211> 29

<212> DNA

25 <213> Artificial Sequence

<220>

<223> single standed oligonucleotide

30 <400> 9

tccatatgaa aagccctgtg gtccaggat

29

<210> 10

35 **<**211> **3**3

<212> DNA

<213> Artificial Sequence

<213>

Artificial Sequence

8

<220>· <223> single stranded oligonucleotide <400> 10 cagtccatat ggtccgccag tgctaccatg gca 33 <210> 11 <211> 31 10 <212> DNA <213> Artificial Sequence <220> 15 <223> single stranded olgonucleotide <400> 11 31 ggaattccat atggaacagg actgcatgtt t 20 <210> 12 <211> 26 <212> DNA <213> Artificial Sequence 25 <220> <223> single stranded oligonucleotide . <400> 12 30 cgggatcctt aacctgattc tgtttc 26 <210> 13 <211> 26 35 <212> DNA

ç

<220> <223> single stranded oligonucleotide

<400> 13

5 cgggatcctt agaccacagt cccttc

26

<210> 14 <211> 23

10 <212> DNA

<213> Artificial Sequence

<220>

<223> single stranded oligonucleotide

15

<400> 14

cgggatcctt aagaggatgc aca